



\ast^s -Modules

Jiaqun Wei

Department of Mathematics, Nanjing Normal University, Nanjing 210097, PR China

Received 13 September 2004

Available online 20 June 2005

Communicated by Efim Zelmanov

Abstract

In this paper we introduce the notion of \ast^s -modules (s denotes *static*) as a generalization of \ast -modules different from \ast^n -modules. The class of \ast^s -modules contains also the class of self-small abelian groups faithfully flat over their endomorphism rings. We study characterizations of \ast^s -modules and extend successfully some results in the theory of \ast -modules. The relations between \ast^s -modules and \ast^n -modules are also considered. Finally we give characterizations of \ast^s -modules with some special properties.

© 2005 Elsevier Inc. All rights reserved.

Keywords: \ast^s -Module; Static modules; \ast -Module

Introduction

The theory of equivalences between module subcategories, originates in the well-known theory of Morita equivalence, had been studied extensively, see [4,8,11,14,19] etc. Modules playing important roles in the theory are the progenerators, quasi-progenerators, tilting modules, \ast -modules and others. In particular, progenerators are just \ast -modules which are generators; quasi-progenerators are just \ast -modules which are Σ -self-generators while tilting modules (of projective dimension ≤ 1) are just \ast -modules which generate all the injective modules.

E-mail address: weijiaqun@njnu.edu.cn.

Recently, the author and the others studied $*^n$ -modules as a generalization of both $*$ -modules and tilting modules of projective dimension $\leq n$ in sense of [12,17,18]. In fact, $*$ -modules are just $*^1$ -modules while tilting modules of projective dimension $\leq n$ are just $*^n$ -modules P such that the subcategory $\text{Pres}^n(P)$ contains all the injective modules and P admits a finitely generated projective resolution.

In this note, we will present another generalization of $*$ -modules which is different from $*^n$ -modules.

Namely, we introduce the notion of $*^s$ -modules, where s denotes *static*, by replacing the subcategory $\text{Gen}(P)$ in the theory of $*$ -modules with the subcategory $\text{Stat}(P)$. Some results on $*$ -modules are successfully extended to our settings (cf. Propositions 2.2, 4.6 and 5.5 etc.). We also compare $*^s$ -modules with $*^n$ -modules. In fact, we show that a $*^s$ -module P is a $*^n$ -module if and only if $\text{Pres}^n(P) \subseteq \text{Stat}(P)$ while a $*^n$ -module P is a $*^s$ -module if and only if $\text{Stat}(P) \subseteq \text{Pres}^n(P)$ (Propositions 3.1 and 3.2). In particular, all $*^2$ -modules are $*^s$ -modules. Interesting, examples of $*^s$ -modules also arise from the theory of abelian groups. Let P be a selfsmall abelian group which is faithfully flat over its endomorphism ring, then P is a $*^s$ -module (Proposition 3.6). We refer to the papers [1,2,10], etc., for the study of such abelian groups. Main characterizations of $*^s$ -modules are given in Section 4. In this section we show that an R -module P with $A = \text{End}_R P$ is a $*^s$ -module if and only if ${}_A \mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ and $\text{Ker } T_P^{i \geq 0} = 0$ (Theorem 4.4). $*^s$ -Modules with some special properties are considered in Section 5. For example, self-small R -modules which are faithfully flat over their endomorphism rings are characterized as $*^s$ -modules flat over their endomorphism rings (Proposition 5.1). We also show the following result (Theorem 5.2). Let P be an R -module. Then P is a $*^s$ -module such that $\text{Stat}(P)$ is a coresolving subcategory, if and only if P is selfsmall and $\text{Stat}(P) = P^\perp$, if and only if $\mathcal{I} \subseteq \text{Stat}(P) \subseteq P^\perp$ and ${}_A \mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$.

1. Preliminaries

Throughout this paper, all rings will be associative with non-zero identity and modules will be left modules. For a ring R , $R\text{-Mod}$ ($\text{Mod-}R$) denotes the category of all left (right) R -modules. By a subcategory, we mean a full subcategory closed under isomorphisms.

From now on, we fix R a ring and P an R -module with $A = \text{End}_R P$. Note that P is also a right A -module. Denote that $H_P = \text{Hom}_R(P, -)$ and $T_P = P \otimes_A -$.

We denote by $P^\perp = \{M \in R\text{-Mod} \mid \text{Ext}_R^i(P, M) = 0 \text{ for all } i \geq 1\}$ and $\text{Ker } T_P^{i \geq n} = \{M \in A\text{-Mod} \mid \text{Tor}_i^A(P, M) = 0 \text{ for all } i \geq n\}$ for a fixed $n \geq 0$ (here we assume that $\text{Tor}_0^A(P, -) = T_P$). Also we denote by $\text{Ad } P$ the class of modules isomorphic to direct sums of copies of the R -module P and by $\text{Add } P$ the class of modules isomorphic to direct summands of modules in $\text{Ad } P$. We denote by ${}_A \mathcal{P}$ the class of all projective A -modules.

An R -module M is n -presented by P (or has P -codominant dimension $\geq n$), for some $n \geq 1$, if there exists an exact sequence $P_n \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow M \rightarrow 0$ with $P_i \in \text{Add } P$ for each i . We denote by $\text{Pres}^n(P)$ the category of all modules with P -codominant dimension $\geq n$. Of course, for every n , we have that $\text{Pres}^{n+1}(P) \subseteq \text{Pres}^n(P)$. Note that $\text{Pres}^2(P)$ and $\text{Pres}^1(P)$ are just the familiar subcategories $\text{Pres}(P)$ and $\text{Gen}(P)$, respectively.

P is said to be (n, t) -quasi-projective (here we assume that $n \geq t \geq 1$) if, for any exact sequence $0 \rightarrow M \rightarrow P_t \rightarrow \cdots \rightarrow P_1 \rightarrow N \rightarrow 0$ with $M \in \text{Pres}^{n-t}(P)$ and $P_i \in \text{Add } P$ for each i , the induced sequence $0 \rightarrow H_P M \rightarrow H_P P_t \rightarrow \cdots \rightarrow H_P P_1 \rightarrow H_P N \rightarrow 0$ is exact [17]. Note that the notions of $(1, 1)$ -quasi-projective, $(2, 1)$ -quasi-projective, $(2, 2)$ -quasi-projective and $(n, 1)$ -quasi-projective respectively are just the notions of Σ -quasi-projective [8,13], w - Σ -quasi-projective [5], semi- Σ -quasi-projective [13] and n -quasi-projective [18], respectively.

P is said to be selfsmall if, for any set X , there is the canonical isomorphism

$$\text{Hom}_R(P, P^{(X)}) \simeq \text{Hom}_R(P, P)^{(X)}.$$

It is well known that (T_P, H_P) is a pair of adjoint functors and there are the following canonical homomorphisms for any R -module M and any A -module N :

$$\begin{aligned} \rho_M : T_P H_P M &\rightarrow M \quad \text{by} \quad p \otimes f \rightarrow f(p); \\ \sigma_N : N &\rightarrow H_P T_P N \quad \text{by} \quad n \rightarrow [p \rightarrow p \otimes n]. \end{aligned}$$

We denote by $\text{Stat}(P)$ the class of P -static modules, i.e., the R -modules M such that ρ_M is an isomorphism. We also denote by $\text{Costat}(P)$ the class of P -costatic modules, i.e., the A -modules N such that σ_N is an isomorphism. Note that the pair of functors (H_P, T_P) defines a basic equivalence between $\text{Stat}(P)$ and $\text{Costat}(P)$. It is also well known that $\text{Stat}(P) \subseteq \text{Pres}(P)$ for any R -module P .

A subcategory is coresolving if it contains all injective objects and is closed under extensions and cokernels of monomorphisms. Dually, a subcategory is resolving if it contains all projective objects and is closed under extensions and kernels of epimorphisms.

Finally, we recall the following characterizations of $*$ -modules. For more results on the theory of $*$ -modules, we refer to the papers [5–7,9,11,15,16], etc.

Theorem 1.1 [5]. *Let P be an R -module with $A = \text{End}_R P$. Then the following statements are equivalent.*

- (1) P is a $*$ -module.
- (2) P is selfsmall, w - Σ -quasi-projective, and $\text{Gen}(P) = \text{Pres}(P)$.
- (3) P is selfsmall and, for any $M \leq P_M$ with $P_M \in \text{Add } P$, $M \in \text{Gen}(P)$ if and only if $\text{Ext}_R^1(P, M) \rightarrow \text{Ext}_R^1(P, P_M)$ is injective canonically.
- (4) P is selfsmall and, for any exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ with $M, N \in \text{Gen}(P)$, the induced sequence $0 \rightarrow H_P L \rightarrow H_P M \rightarrow H_P N \rightarrow 0$ is exact if and only if $L \in \text{Gen}(P)$.

2. Basic properties on $*^s$ -modules

Note that, for P a $*$ -module, the subcategory $\text{Stat}(P)$ equals $\text{Gen}(P)$ [5], so it is interesting to consider what happens if we replace the subcategory $\text{Gen}(P)$ in the theory of $*$ -modules with the subcategory $\text{Stat}(P)$. This leads the following definition.

Definition 2.1. An R -module P is said to be a $*^s$ -module provided that P is selfsmall and that any exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ with $M, N \in \text{Stat}(P)$ remains exact after applying the functor H_P if and only if $L \in \text{Stat}(P)$.

It is easy to see that $*^s$ -modules with $\text{Stat}(P) = \text{Gen}(P)$ are just $*$ -modules (cf. Theorem 1.1). Hence the notion of $*^s$ -modules is a generalization of $*$ -modules.

For a $*^s$ -module P , the subcategory $\text{Stat}(P)$ has the following properties.

Proposition 2.2. *Let P be a $*^s$ -module. Then*

- (1) *The functor H_P preserves short exact sequence s in $\text{Stat}(P)$.*
- (2) *For any $M \in \text{Stat}(P)$, there is an infinite exact sequence $\cdots \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow M \rightarrow 0$ which remains exact after applying the functor H_P , where $P_i \in \text{Add } P$ for each i .*
- (3) *For any exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ which is also exact after applying the functor H_P , if two of its terms are in $\text{Stat}(P)$, then so is the third one.*

Proof. (1) Follows from the definition of $*^s$ -modules.

(2) Note that $\text{Stat}(P) \subseteq \text{Pres}(P)$, so, for any $M \in \text{Stat}(P)$, we have an exact sequence $0 \rightarrow M_1 \rightarrow P_1 \xrightarrow{f_1} M \rightarrow 0$ with $P_1 = P^{(H_P M)} \in \text{Add } P$ and f_1 the evaluation map. Since the sequence is clearly exact after applying the functor H_P and $P_1, M \in \text{Stat}(P)$, we obtain that $M_1 \in \text{Stat}(P)$. By repeating the process to M_1 , and so on, we finally obtain the desired exact sequence.

(3) If $M, N \in \text{Stat}(P)$, then $L \in \text{Stat}(P)$ by assumptions and Definition 2.1.

Now let $L \in \text{Stat}(P)$. By applying the functor $T_P H_P$ to the sequence in (3), we have the following exact commutative diagram by assumptions.

$$\begin{array}{ccccccc}
 T_P H_P L & \longrightarrow & T_P H_P M & \longrightarrow & T_P H_P N & \longrightarrow & 0 \\
 \downarrow \rho_L & & \downarrow \rho_M & & \downarrow \rho_N & & \\
 0 & \longrightarrow & L & \longrightarrow & M & \longrightarrow & N \longrightarrow 0
 \end{array}$$

Note that ρ_L is an isomorphism since $L \in \text{Stat}(P)$. It follows easily that ρ_M is an isomorphism if and only if ρ_N is an isomorphism. Hence $M \in \text{Stat}(P)$ if and only if $N \in \text{Stat}(P)$. \square

We note that the converse of Proposition 2.2(2) holds whenever P is selfsmall [19].

The following proposition gives some properties of the subcategory $\text{Costat}(P)$ for P a $*^s$ -module.

Proposition 2.3. *Let P be a $*^s$ -module with $A = \text{End}_R P$. Then*

- (1) ${}_A \mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$. *In particular, the functor T_P preserves short exact sequences in $\text{Costat}(P)$.*
- (2) $\text{Costat}(P)$ *is a resolving subcategory.*

Proof. (1) Since P is selfsmall, we easily obtain that ${}_A\mathcal{P} \subseteq \text{Costat}(P)$. Now for any $N \in \text{Costat}(P)$, we have that $T_P N \in \text{Stat}(P)$, and hence there is an infinite exact sequence $\cdots \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow T_P N \rightarrow 0$ which remains exact after applying the functor H_P , where $P_i \in \text{Add } P$ for each i , by Proposition 2.2(2). Then, by applying the functor H_P to the sequence, we obtain an exact sequence $\cdots \rightarrow H_P P_n (\simeq A_n) \rightarrow \cdots \rightarrow H_P P_1 (\simeq A_1) \rightarrow H_P T_P N (\simeq N) \rightarrow 0$ with $A_i \in {}_A\mathcal{P}$. Since it is clearly exact after applying the functor T_P , we obtain that $\text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ by dimension shifting.

(2) $\text{Costat}(P)$ contains all projective A -modules by (1). Now for any exact sequence $0 \rightarrow K \rightarrow M \rightarrow N \rightarrow 0$ with $N \in \text{Costat}(P)$, we obtain the following exact commutative diagram by applying the functor $H_P T_P$ since $\text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ by (1).

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K & \longrightarrow & M & \longrightarrow & N & \longrightarrow & 0 \\ & & \downarrow \sigma_K & & \downarrow \sigma_M & & \downarrow \sigma_N & & \\ 0 & \longrightarrow & H_P T_P K & \longrightarrow & H_P T_P M & \longrightarrow & H_P T_P N & \longrightarrow & \end{array}$$

Note that σ_N is an isomorphism since $N \in \text{Costat}(P)$, we have that σ_K is an isomorphism if and only if σ_M is an isomorphism. That is, $K \in \text{Costat}(P)$ if and only if $M \in \text{Costat}(P)$. It follows that $\text{Costat}(P)$ is also closed under kernels of epimorphisms and under extensions. Thus, $\text{Costat}(P)$ is a resolving subcategory. \square

3. Relations between $*^s$ -modules and $*^n$ -modules

The notion of $*^n$ -modules, introduced in [17,18] by the author and the others, is also a generalization of $*$ -modules. It is natural to consider the relations between $*^s$ -modules and $*^n$ -modules. Recall that P is said to be a $*^n$ -module if P is selfsmall, $(n+1)$ -quasi-projective and $\text{Pres}^n(P) = \text{Pres}^{n+1}(P)$ [18]. Note that $*$ -modules are exactly $*^1$ -modules (cf. Theorem 1.1).

The following is the first result in this direction.

Proposition 3.1. *Let P be a $*^s$ -module. Then P is a $*^n$ -module if and only if $\text{Pres}^n(P) \subseteq \text{Stat}(P)$. In particular, P is a $*^n$ -module provided that P is n -quasi-projective with $n \geq 2$.*

Proof. If P is a $*^n$ -module, then we clearly have that $\text{Pres}^n(P) \subseteq \text{Stat}(P)$ by [18]. Now assume that $\text{Pres}^n(P) \subseteq \text{Stat}(P)$. By Proposition 2.2, we have that $\text{Stat}(P) \subseteq \text{Pres}^n(P)$. It follows that $\text{Stat}(P) = \text{Pres}^n(P)$. Hence we obtain that P is selfsmall and any exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ with $M, N \in \text{Pres}^n(P)$ remains exact after applying the functor H_P if and only if $L \in \text{Pres}^n(P)$ by Definition 2.1. Therefore, P is also a $*^n$ -module by [18].

If P is n -quasi-projective, then $\text{Pres}^n(P) \subseteq \text{Stat}(P)$ by [17]. Hence P is a $*^n$ -module by the first part. \square

Similarly, we have the following result.

Proposition 3.2. *Let P be a $*^n$ -module. Then P is a $*^s$ -module if and only if $\text{Stat}(P) \subseteq \text{Pres}^n(P)$.*

Proof. The necessity has been proved in the last proposition. Now assume that $\text{Stat}(P) \subseteq \text{Pres}^n(P)$, then we obtain that $\text{Pres}^n(P) = \text{Stat}(P)$ since $\text{Pres}^n(P) \subseteq \text{Stat}(P)$ due to that P is a $*^n$ -module. Hence, by [18] and the definition of $*^s$ -modules, we have that P is a $*^s$ -module. \square

A corollary of the last proposition shows that all $*^2$ -modules are $*^s$ -modules.

Corollary 3.3. *Let P be a $*^2$ -module. Then P is a $*^s$ -module.*

As to cases $n \geq 3$, we will see that $*^n$ -modules need not be $*^s$ -modules from the following example.

Example 3.4. A $*^n$ -module ($n \geq 3$) need not be a $*^s$ -module.

Proof. In fact, let R denote the path algebra defined by the quiver $1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3$ with the relation $\alpha\beta = 0$. Then, for this algebra, we have a projective module $P = \begin{smallmatrix} 2 \\ 1 \end{smallmatrix} \oplus \begin{smallmatrix} 3 \\ 2 \end{smallmatrix}$. Since $\text{Pres}^3(P) = \text{Add } P$, we see that P is a $*^3$ -module [18]. Note that the simple module $S(3) \in \text{Pres}^2(P)$ and that $S(3) \notin \text{Pres}^3(P)$, so we obtain that $\text{Pres}^3(P) \subsetneq \text{Pres}^2(P)$ and hence P is not a $*^2$ -module. Since P is projective, we have that P cannot be a $*^s$ -module. Otherwise, P must be a $*^2$ -module by Proposition 3.1 since projective modules are clearly 2-quasi-projective, a contradiction. \square

Using the notion of $*^s$ -modules, progenerators and quasi-progenerators can be characterized as follows.

Proposition 3.5. *Let P be an R -module.*

- (1) *P is a progenerator if and only if P is a $*^s$ -module and a generator.*
- (2) *P is a quasi-progenerator if and only if P is a $*^s$ -module and a Σ -self-generator.*

Proof. The necessity is clear. Now if P is a Σ -self-generator, then $\text{Gen}(P) = \text{Stat}(P)$. Since P is also a $*^s$ -module, P is a $*^s$ -module by Proposition 3.1. An R -module which is a $*^s$ -module and a generator (a Σ -self-generator, respectively) is clearly a progenerator (a quasi-progenerator, respectively) [5], so the sufficient parts hold. \square

It is interesting to know if there are other $*^s$ -modules which are not $*^n$ -modules for any n . The following result will give an answer to this question.

Proposition 3.6. *Let P be a selfsmall R -module with $A = \text{End}_R P$. If P_A is faithfully flat, then P is a $*^s$ -module.*

Proof. Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be an exact sequence with $M, N \in \text{Stat}(P)$. Assume first that the sequence is exact after applying the functor H_P , i.e., we have the induced exact sequence $0 \rightarrow H_P L \rightarrow H_P M \rightarrow H_P N \rightarrow 0$. Since P_A is flat by assumptions, after applying the functor T_P we obtain that the sequence $0 \rightarrow T_P H_P L \rightarrow T_P H_P M \rightarrow T_P H_P N \rightarrow 0$ is exact. Note that $M, N \in \text{Stat}(P)$, so ρ_M and ρ_N are isomorphisms. It follows that ρ_L is an isomorphism and $L \in \text{Stat}(P)$.

Now suppose that $L \in \text{Stat}(P)$. After applying the functor H_P , we obtain an induced exact sequence $0 \rightarrow H_P L \rightarrow H_P M \rightarrow H_P N \rightarrow D \rightarrow 0$ for some D . Since also P_A is flat, we have that the sequence $0 \rightarrow T_P H_P L \rightarrow T_P H_P M \rightarrow T_P H_P N \rightarrow T_P D \rightarrow 0$ is exact, by again applying the functor T_P . It is easy to see that $T_P D = 0$. Since P_A is faithfully flat, we then obtain that $D = 0$ [3]. Hence $0 \rightarrow H_P L \rightarrow H_P M \rightarrow H_P N \rightarrow 0$ is exact. By Definition 2.1, we have that P is a $*^s$ -module. \square

We note that selfsmall abelian groups (i.e., \mathbf{Z} -modules) which are faithfully flat over their endomorphism rings play important roles in the studies of the theory of abelian groups (see for instance [1,2,10] etc.). The following is one example of such abelian groups. We refer to the paper [10] and the references cited there for more examples. Note that the example also shows that, in contrast to $*$ -modules which are always finitely generated, $*^s$ -modules are not finitely generated in general.

Example 3.7. Let P be a torsion-free abelian group of rank 2 with $\text{End } P = \mathbf{Z}_p$. Then P is selfsmall abelian group and P is faithfully flat over its endomorphism ring \mathbf{Z}_p . In the case, P is a $*^s$ -module which is not a $*^n$ -module for any n .

Proof. P is selfsmall abelian group and P is faithfully flat over its endomorphism ring $A = \mathbf{Z}_p$, as shown in [1, Example 3.3]. To see that P is never a $*^n$ -module for any n , it is sufficient to show that $\text{Costat}(P) \neq \text{Ker } T_P^{i \geq 1} = A\text{-Mod}$ by [18]. However, this is followed from the fact that, in the case, $\text{Costat}(P)$ is not a torsion-free class by [1, Example 3.3 and Theorem 2.3]. \square

4. Characterizations of $*^s$ -modules

Suggested by Proposition 3.6, we have the following more general result.

Proposition 4.1. Let P be an R -module with $A = \text{End}_R P$. Assume that ${}_A \mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ and $\text{Ker } T_P^{i \geq 0} = 0$. Then P is a $*^s$ -module.

Proof. It is easy to see that ${}_A \mathcal{P} \subseteq \text{Costat}(P)$ implies that P is selfsmall.

Now let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be an exact sequence with $M, N \in \text{Stat}(P)$. Assume first that the sequence is exact after applying the functor H_P , i.e., we have the induced exact sequence $0 \rightarrow H_P L \rightarrow H_P M \rightarrow H_P N \rightarrow 0$. Since $H_P N \in \text{Ker } T_P^{i \geq 1}$, we obtain $L \in \text{Stat}(P)$ as in the proof of Proposition 3.6.

Now suppose that $L \in \text{Stat}(P)$. By applying the functor H_P , we obtain an induced exact sequences $0 \rightarrow H_P L \rightarrow H_P M \rightarrow X \rightarrow 0$ and $0 \rightarrow X \xrightarrow{f} H_P N \rightarrow Y \rightarrow 0$ for some

X, Y . After applying the functor T_P to the first sequence, we then have the following exact commutative diagram.

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & \mathrm{Tor}_1^A(P, X) & \longrightarrow & T_P H_P L & \longrightarrow & T_P H_P M & \longrightarrow & T_P X & \longrightarrow & 0 \\
 & & & & \downarrow \rho_L & & \downarrow \rho_M & & \downarrow h & & \\
 0 & \longrightarrow & L & \longrightarrow & M & \longrightarrow & N & \longrightarrow & 0
 \end{array}$$

Note that $H_P L, H_P M \in \mathrm{Costat}(P) \subseteq \mathrm{Ker} T_P^{i \geq 1}$, so we have that $\mathrm{Tor}_i^A(P, X) = 0$ for all $i \geq 2$ by dimension shifting. Moreover, ρ_L and ρ_M are isomorphisms, since $L, M \in \mathrm{Stat}(P)$. It follows from the last diagram that $\mathrm{Tor}_1^A(P, X) = 0$ and $h = (T_P f) \rho_N$ is an isomorphism. Now we have that $X \in \mathrm{Ker} T_P^{i \geq 1}$ and that $T_P f$ is an isomorphism since ρ_N is an isomorphism. Note also that, by applying the functor T_P to the sequence $0 \rightarrow X \xrightarrow{f} H_P N \rightarrow Y \rightarrow 0$, we have an induced exact sequence

$$0 \rightarrow \mathrm{Tor}_1^A(P, Y) \rightarrow T_P X \xrightarrow{T_P f} T_P H_P N \rightarrow T_P Y \rightarrow 0$$

and that $\mathrm{Tor}_i^A(P, Y) = 0$ for all $i \geq 2$, since $H_P N \in \mathrm{Costat}(P) \subseteq \mathrm{Ker} T_P^{i \geq 1}$. Hence we obtain that

$$\mathrm{Tor}_1^A(P, Y) = 0 = T_P Y$$

by arguments above. It follows that $Y \in \mathrm{Ker} T_P^{i \geq 0}$. Then $Y = 0$ by assumptions and hence $X \simeq H_P N$ canonically. Therefore, we deduce that the functor H_P preserves the exactness of the exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ in $\mathrm{Stat}(P)$.

Finally, we conclude that P is a $*^s$ -module. \square

In fact, the assumptions in the last result also characterize $*^s$ -modules. To see this, we need the following lemma.

Lemma 4.2. *Let P be a selfsmall R -module with $A = \mathrm{End}_R P$. Then $\Omega^2(\mathrm{Ker} T_P^{i \geq 1}) \subseteq \mathrm{Costat}(P)$, where Ω^2 denotes the second syzygy module.*

Proof. For any $N \in \Omega^2(\mathrm{Ker} T_P^{i \geq 1})$, we have an exact sequence $0 \rightarrow N \rightarrow A_2 \rightarrow A_1 \rightarrow M \rightarrow 0$ with $A_1, A_2 \in {}_A \mathcal{P}$ and $M \in \mathrm{Ker} T_P^{i \geq 1}$. Then we have an induced exact sequence $0 \rightarrow T_P N \rightarrow T_P A_2 \rightarrow T_P A_1 \rightarrow T_P M \rightarrow 0$ by applying the functor T_P . Now after applying the functor H_P to the last sequence, we obtain an induced exact sequence $0 \rightarrow H_P T_P N \rightarrow H_P T_P A_2 \rightarrow H_P T_P A_1$. Note that σ_{A_2} and σ_{A_1} are isomorphisms since P is selfsmall, so we deduce that σ_N is also an isomorphism. Hence $N \in \mathrm{Costat}(P)$. \square

Let P be a $*^s$ -module. We have seen that ${}_A \mathcal{P} \subseteq \mathrm{Costat}(P) \subseteq \mathrm{Ker} T_P^{i \geq 1}$ in Proposition 2.3. Now we will show that $\mathrm{Ker} T_P^{i \geq 0} = 0$ in this case.

Proposition 4.3. *Let P be a $*^s$ -module. Then $\mathrm{Ker} T_P^{i \geq 0} = 0$.*

Proof. For any $M \in \text{Ker } T_P^{i \geq 0}$, we have an exact sequence $0 \rightarrow N \rightarrow A_2 \rightarrow A_1 \rightarrow M \rightarrow 0$ with $A_1, A_2 \in {}_A\mathcal{P}$. Then, by applying the functor T_P to the sequence, we obtain an induced exact sequence $0 \rightarrow T_P N \rightarrow T_P A_2 \rightarrow T_P A_1 \rightarrow 0$. Note that $T_P N \in \text{Stat}(P)$ since $N \in \text{Costat}(P)$ by Lemma 4.2, so, after applying the functor H_P , we obtain that the induced sequence $0 \rightarrow H_P T_P N \rightarrow H_P T_P A_2 \rightarrow H_P T_P A_1 \rightarrow 0$ is exact by Proposition 2.2. It follows that $M = \text{Coker}(A_2 \rightarrow A_1) \simeq \text{Coker}(H_P T_P A_2 \rightarrow H_P T_P A_1) = 0$. \square

Combining Propositions 2.3, 4.1 and 4.3, we obtain the following characterization of $*^s$ -modules.

Theorem 4.4. *Let P be an R -module with $A = \text{End}_R P$. Then P is a $*^s$ -module if and only if ${}_A\mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ and $\text{Ker } T_P^{i \geq 0} = 0$.*

Specially, we have the following result.

Proposition 4.5. *Let P be an R -module with $A = \text{End}_R P$. Assume that $\text{Costat}(P) = \text{Ker } T_P^{i \geq 1}$, then P is a $*^s$ -module.*

Proof. Under assumptions we clearly have that ${}_A\mathcal{P} \subseteq \text{Costat}(P)$. Moreover, for any $M \in \text{Ker } T_P^{i \geq 0}$, we obtain that $M \in \text{Costat}(P) = \text{Ker } T_P^{i \geq 1}$. Hence $M \simeq H_P T_P M = 0$. By Theorem 4.4, we see that P is a $*^s$ -module. \square

We can now give a characterization of $*^s$ -modules which is similar to Theorem 1.1(3).

Proposition 4.6. *Let P be a selfsmall R -module. The following are equivalent.*

- (1) P is a $*^s$ -module.
- (2) For any exact sequence $0 \rightarrow L \rightarrow P_M \rightarrow M \rightarrow 0$ with $P_M \in \text{Add } P$ and $M \in \text{Stat}(P)$, $L \in \text{Stat}(P)$ if and only if the functor H_P preserves the exactness of the sequence.

Proof. (1) \Rightarrow (2) is followed from the definition of $*^s$ -module.

(2) \Rightarrow (1). As proved in Proposition 2.2(2), we can obtain that, for any $M \in \text{Stat}(P)$, there is an infinite exact sequence $\cdots \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow M \rightarrow 0$ which remains exact after applying the functor H_P , where $P_i \in \text{Add } P$ for each i . Consequently, we have that ${}_A\mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ as in the proof of Proposition 2.3(1). Finally, the same proof as in Proposition 4.3 yields that $\text{Ker } T_P^{i \geq 0} = 0$. Thus we obtain that P is a $*^s$ -module by Theorem 4.4. \square

5. $*^s$ -Modules with special properties

This section is devoted to the study of $*^s$ -modules P such that the subcategory $\text{Stat}(P)$ has some special properties.

We have seen that selfsmall R -modules which are faithfully flat over their endomorphism rings are $*^s$ -modules before. The following result shows that they are just $*^s$ -modules which are flat over their endomorphism rings.

Proposition 5.1. *Let P be a selfsmall R -module with $A = \text{End}_R P$. The following are equivalent.*

- (1) P_A is faithfully flat.
- (2) P is a $*^s$ -module and P_A is flat.
- (3) P is a $*^s$ -module and $\text{Stat}(P)$ is closed under kernels of homomorphisms.

Proof. (1) \Rightarrow (2). By Proposition 3.6.

(2) \Leftrightarrow (3). It is well known that P_A is flat if and only if $\text{Stat}(P)$ is closed under kernels of homomorphisms (see for instance [19]).

(2) \Rightarrow (1). Since P_A is flat and P is a $*^s$ -module, we have that $\text{Ker } T_P = \text{Ker } T_P^{i \geq 0} = 0$ by Proposition 4.3. Hence P_A is faithfully flat [3]. \square

For a $*^s$ -module P , we know that $\text{Costat}(P)$ is a resolving subcategory from Proposition 2.3(2). Dually, we can consider when $\text{Stat}(P)$ is a coresolving subcategory. The following gives some characterizations of this case. We note that the class of such $*^s$ -modules contains all tilting modules of projective dimension ≤ 1 .

Theorem 5.2. *Let P be an R -module and \mathcal{I} be the class of all injective R -modules. The following are equivalent.*

- (1) P is a $*^s$ -module such that $\text{Stat}(P)$ is a coresolving subcategory.
- (2) P is selfsmall and $\text{Stat}(P) = P^\perp$.
- (3) P is selfsmall and $\text{Stat}(P) \subseteq P^\perp \subseteq \text{Gen}(P)$.
- (4) $\mathcal{I} \subseteq \text{Stat}(P) \subseteq P^\perp$ and ${}_A\mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$.

Proof. (1) \Rightarrow (2). Let $S \in \text{Stat}(P)$ and $0 \rightarrow S \rightarrow I_1 \rightarrow S_1 \rightarrow 0$ be an exact sequence with $I_1 \in \mathcal{I}$. Since $\text{Stat}(P)$ is a coresolving subcategory, $\mathcal{I} \subseteq \text{Stat}(P)$ and $\text{Stat}(P)$ is closed under cokernels of monomorphisms. Hence we have that $S_1 \in \text{Stat}(P)$. It follows from Proposition 2.2 that the sequence remains exact after applying the functor H_P . Then we obtain that $\text{Ext}_R^1(P, S) = 0$ for any $S \in \text{Stat}(P)$. Therefore, $\text{Stat}(P) \subseteq P^\perp$ since $\text{Stat}(P)$ is a coresolving subcategory.

On the other hand, we may consider the exact sequence $0 \rightarrow S \rightarrow I_1 \rightarrow I_2 \rightarrow X \rightarrow 0$ with $I_1, I_2 \in \mathcal{I}$, for any $S \in P^\perp$. It clearly remains exact after applying the functor H_P . Hence we have an induced exact sequence $0 \rightarrow H_P S \rightarrow H_P I_1 \rightarrow H_P I_2 \rightarrow H_P X \rightarrow 0$. By applying the functor T_P to the sequence, we obtain that $X = \text{Coker}(I_1 \rightarrow I_2) \simeq \text{Coker}(T_P H_P I_1 \rightarrow T_P H_P I_2) = T_P H_P X$ canonically since $\mathcal{I} \subseteq \text{Stat}(P)$. That is, $X \in \text{Stat}(P)$. Since P is a $*^s$ -module, we see that $\text{Im}(I_1 \rightarrow I_2) \in \text{Stat}(P)$ and similarly $S \in \text{Stat}(P)$. Hence we have also $P^\perp \subseteq \text{Stat}(P)$.

(2) \Rightarrow (3) is obvious.

(3) \Rightarrow (4). By assumptions we have that $\text{Add } P \subseteq P^\perp$. Now for any $M \in P^\perp$, we have an exact sequence $0 \rightarrow M_1 \rightarrow P_1 \xrightarrow{f_1} M \rightarrow 0$ with $P_1 \in P^{(H_P M)}$ and f_1 the evaluation map, since $P^\perp \subseteq \text{Gen}(P)$. Note the sequence clearly stays exact after applying the functor H_P and since $P_1, M \in P^\perp$, we obtain that $M_1 \in P^\perp$ too. By repeating the process to M_1 , and so on, we finally obtain an infinite exact sequence $\cdots \rightarrow P_n \rightarrow \cdots \rightarrow P_1 \rightarrow M \rightarrow 0$, with $P_i \in \text{Add } P$ for each i , such that the sequence remains exact after applying the functor H_P . It follows that $M \in \text{Stat}(P)$ by [19]. Hence we deduce that $\text{Stat}(P) = P^\perp$. Moreover, by an argument similar to the proof of Proposition 2.3(1), we obtain that ${}_A \mathcal{P} \subseteq \text{Costat}(P) \subseteq \text{Ker } T_P^{i \geq 1}$ since P is also selfsmall.

(4) \Rightarrow (1). Since $\mathcal{I} \subseteq \text{Stat}(P) \subseteq P^\perp$, we easily check that $\text{Stat}(P)$ is a coresolving subcategory by a dual proof of Proposition 2.3(2). Now it remains to show that P is a $*^s$ -module. By Theorem 4.4, we need only to prove that $\text{Ker } T_P^{i \geq 0} = 0$. This will be proved similarly as in Proposition 4.3. Namely, let $M \in \text{Ker } T_P^{i \geq 0}$ and take an exact sequence $0 \rightarrow N \rightarrow A_2 \rightarrow A_1 \rightarrow M \rightarrow 0$ with $A_2, A_1 \in {}_A \mathcal{P}$. Then we have an induced exact sequence $0 \rightarrow T_P N \rightarrow T_P A_2 \rightarrow T_P A_1 \rightarrow 0$ by applying the functor T_P . Note that $N \in \text{Costat}(P)$ by Lemma 4.2, so that $T_P N \in \text{Stat}(P) \subseteq P^\perp$. It follows that there is an induced exact sequence $0 \rightarrow H_P T_P N \rightarrow H_P T_P A_2 \rightarrow H_P T_P A_1 \rightarrow 0$ by applying the functor H_P . Hence we obtain that $M = \text{Coker}(A_2 \rightarrow A_1) \simeq \text{Coker}(H_P T_P A_2 \rightarrow H_P T_P A_1) = 0$. \square

In particular, if P is a small $*^s$ -module and $\text{Stat}(P)$ is coresolving and closed under homomorphic images, then P^\perp is a 1-tilting class.

Clearly, for P a $*^s$ -module, $\text{Stat}(P)$ cannot be resolving except that P is a progenitor. However, it may happen that $\text{Stat}(P)$ is closed under kernels of epimorphisms or extensions.

Recall that an R -module P is Σ -direct-projective if any epimorphism $f : L \rightarrow N$ with $L, N \in \text{Add } P$ splits (see for instance [19]).

Proposition 5.3. *Let P be a $*^s$ -module. The following are equivalent.*

- (1) $\text{Stat}(P)$ is closed under kernels of epimorphisms.
- (2) P is Σ -direct-projective.
- (3) $\text{Ker } T_P = 0$.

Proof. (1) \Rightarrow (2). For any epimorphism $f : P_1 \rightarrow P_0$ with $P_0, P_1 \in \text{Add } P$, let L be the kernel of f , then $L \in \text{Stat}(P)$ by assumption. Since P is a $*^s$ -module, we have an induced exact sequence $0 \rightarrow H_P L \rightarrow H_P P_1 \rightarrow H_P P_0 \rightarrow 0$ after applying the functor H_P by Proposition 2.2(1). Note that the sequence splits since $H_P P_0$ is projective, so we obtain that $f \simeq T_P H_P f$ splits too. It follows that P is Σ -direct-projective.

(2) \Rightarrow (3). Let $M \in \text{Ker } T_P$ and take an exact sequence $A_1 \xrightarrow{f} A_0 \rightarrow M \rightarrow 0$ with $A_0, A_1 \in {}_A \mathcal{P}$. Then we have an induced exact sequence $T_P A_1 \xrightarrow{T_P f} T_P A_0 \rightarrow 0$ by applying the functor T_P . Since P is Σ -direct-projective, $T_P f$ splits. It follows that the induced sequence $H_P T_P A_1 \xrightarrow{H_P T_P f} H_P T_P A_0 \rightarrow 0$ is exact. Hence we have that $M = \text{Coker } f \simeq \text{Coker } H_P T_P f = 0$.

(3) \Rightarrow (1). For any exact sequence

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0 \quad (*)$$

with $M, N \in \text{Stat}(P)$, we have an induced exact sequence $0 \rightarrow H_P L \rightarrow H_P M \rightarrow H_P N \rightarrow X \rightarrow 0$ by applying the functor H_P . By applying the functor T_P to the last sequence we also have an induced exact sequence $T_P H_P M \rightarrow T_P H_P N \rightarrow T_P X \rightarrow 0$. Then, $T_P X = \text{Coker}(T_P H_P M \rightarrow T_P H_P N) \simeq \text{Coker}(M \rightarrow N) = 0$. It follows that $X = 0$ by assumptions. Hence the exact sequence $(*)$ remains exact after applying the functor H_P . Since P is a $*^s$ -module, we conclude that $L \in \text{Stat}(P)$ and $\text{Stat}(P)$ is closed under kernels of epimorphisms. \square

As a corollary, we have the following result.

Corollary 5.4.

- (1) P is a quasi-progenerator if and only if P is a Σ -direct-projective $*^s$ -module.
- (2) P is a 2-quasi-progenerator if and only if P is a Σ -direct-projective $*^2$ -module.

Proof. (1) The necessary part is obvious. As to the sufficient part, note that $\text{Gen}(P) = \text{Stat}(P)$ since P is a $*^s$ -module, so we have that $L \in \text{Gen}(P)$, for any exact sequence $0 \rightarrow L \rightarrow P' \rightarrow N \rightarrow 0$ with $P' \in \text{Add } P$, by assumptions and Proposition 5.3. In another word, P is a Σ -self-generator. Now the conclusion follows from Proposition 3.5.

(2) Similarly, we need only to show that, if P is a Σ -direct-projective $*^2$ -module, then P is a 2-quasi-progenerator. Note that $\text{Pres}(P) = \text{Stat}(P)$ since P is a $*^2$ -module. Now consider any exact sequence $0 \rightarrow L \rightarrow P_2 \rightarrow P_1 \rightarrow N \rightarrow 0$ with $P_1, P_2 \in \text{Add } P$ and let $M = \text{Ker}(P_1 \rightarrow N)$. By Proposition 5.3, we obtain that $M \in \text{Pres}(P)$, and consequently $L \in \text{Pres}(P)$. It follows that the sequence remains exact after applying the functor H_P by Proposition 2.2(1). Hence P is (2, 2)-quasi-projective and, consequently, is a 2-quasi-progenerator by [17]. \square

We end with the following result, which deals with when the subcategory $\text{Stat}(P)$ is closed under extensions. The proof is similar to that of the corresponding result on $*^s$ -modules [5], so we omit it here.

Proposition 5.5. *Let P be a $*^s$ -module. Then $\text{Stat}(P)$ is closed under extensions if and only if $\text{Ext}_R^1(P, M) = 0$ for any $M \in \text{Stat}(P)$.*

References

- [1] U. Albrecht, The construction of A -solvable abelian groups, Czech. Math. J. 44 (119) (1994) 413–430.
- [2] U. Albrecht, Baer's Lemma and Fuchs' Problem 84a, Trans. Amer. Math. Soc. 293 (2) (1986) 565–582.
- [3] F.D. Anderson, K.R. Fuller, Rings and Categories of Modules, first ed., Springer, New York, 1974.
- [4] S. Brenner, M.C.R. Butler, Generalizations of the Bernstein–Gelfand–Ponomarev reflection functors, in: Lecture Notes in Math., vol. 832, Springer, 1980, pp. 103–170.

- [5] R. Colpi, Some remarks on equivalences between categories of modules, *Comm. Algebra* 18 (1990) 1935–1951.
- [6] R. Colpi, Tilting modules and $*$ -modules, *Comm. Algebra* 21 (4) (1993) 1095–1102.
- [7] R. Colpi, C. Menini, On the structure of $*$ -modules, *J. Algebra* 158 (2) (1993) 400–419.
- [8] K.R. Fuller, Density and equivalence, *J. Algebra* 29 (1974) 528–550.
- [9] K.R. Fuller, $*$ -Modules over ring extensions, *Comm. Algebra* 29 (9) (1997) 2839–2860.
- [10] T.G. Faticoni, P. Gorters, Examples of torsion-free groups flat as modules over their endomorphism rings, *Comm. Algebra* 19 (1) (1991) 1–27.
- [11] C. Menini, A. Orsatti, Representable equivalences between categories of modules and applications, *Rend. Sem. Mat. Univ. Padova* 82 (1989) 203–231.
- [12] Y. Miyashita, Tilting modules of finite projective dimension, *Math. Z.* 193 (1986) 113–146.
- [13] M. Sato, Fuller’s Theorem on equivalences, *J. Algebra* 52 (1978) 174–184.
- [14] M. Sato, On equivalences between module categories, *J. Algebra* 59 (2) (1979) 412–420.
- [15] J. Trlifaj, $*$ -Modules are finitely generated, *J. Algebra* 169 (1994) 392–398.
- [16] J. Trlifaj, Dimension estimates for representable equivalences of module categories, *J. Algebra* 193 (1997) 660–676.
- [17] J. Wei, (n, t) -Quasi-projective and equivalences, *Comm. Algebra*, in press.
- [18] J. Wei, Z. Huang, W. Tong, J. Huang, Tilting modules of finite projective dimension and a generalization of $*$ -modules, *J. Algebra* 168 (2) (2003) 404–418.
- [19] R. Wisbauer, Static modules and equivalences, in: *Interactions between Ring Theory and Representations of Algebras*, Murcia, in: *Lecture Notes in Pure and Appl. Math.*, vol. 210, Dekker, New York, 2000, pp. 423–449.